



Exploring the case for research on incorporating psychedelics within interventions for borderline personality disorder[☆]



Richard J. Zeifman^{a,*}, Anne C. Wagner^{a,b}

^a Department of Psychology, Ryerson University, 350 Victoria St, Toronto, M5B 2K3, Canada

^b Remedy, 703 Bloor Street West, Toronto, M6G 1L5, Canada

ARTICLE INFO

Keywords:

Borderline personality disorder
Psychedelics
Psychotherapy
Pharmacotherapy
Mechanisms of change

ABSTRACT

Borderline Personality Disorder (BPD) is a severe psychiatric disorder characterized by behavioral dysregulation, emotion dysregulation, disturbances in self-identity, and social functioning. Despite the existence of evidence-based psychotherapeutic interventions for BPD, these interventions have important limitations (e.g., limited treatment efficacy). Furthermore, little evidence exists for the efficaciousness of pharmacological interventions for BPD. Thus, a strong need for improving current interventions for BPD exists. Although research incorporating psychedelics within interventions for a wide range of psychiatric disorders has shown promise, research has not yet explored the utility of incorporating psychedelics within interventions for BPD. Therefore, this paper reviews the impact of psychedelics on treatment targets in interventions for BPD (i.e., behavioral dysregulation, emotion dysregulation, disturbances in self-identity, and social functioning), as well as purported overlapping mechanisms of change between interventions for BPD and psychedelics (i.e., emotion dysregulation, mindfulness, and self-compassion). Finally, safety concerns, clinical recommendations, and proposed next research steps related to the administration of psychedelics to individuals with BPD are discussed. This paper aims to explore the case for conducting research on, and the potential clinical utility surrounding, incorporating psychedelics within interventions for BPD.

Borderline personality disorder (BPD) is a severe psychiatric disorder characterized by behavioral dysregulation, emotion dysregulation, disturbances in self-identity, and difficulties with social functioning (American Psychiatric Association [APA], 2013). BPD has a prevalence rate of approximately 1.5% in the general population and is especially prevalent within clinical populations (9.3%–46.3%; Torgersen, 2012). Moreover, BPD is associated with high rates of suicide (Temes, Frankenburg, Fitzmaurice, & Zanarini, 2019), intensive use of healthcare resources (Caillhol, Thalamas, Garrido, Birmes, & Lapeyre-Mestre, 2015; Zanarini, Frankenburg, Hennen, & Silk, 2004), and, due to healthcare utilization and reduced involvement in the workforce, creates significant economic burden on society (Van Asselt, Dirksen, Arntz, & Severens, 2007). Therefore, efficacious interventions for BPD are of great importance.

1. The current state of interventions for borderline personality disorder

1.1. Psychotherapeutic interventions for borderline personality disorder

Psychotherapeutic interventions are currently considered the first-line treatment for BPD (Paris, 2009). A number of psychotherapeutic interventions have been developed specifically for BPD, including dialectical behavior therapy (DBT; Linehan, 1993), transference-focused psychotherapy (TFP; Clarkin, Yeomans, & Kernberg, 1999), and mentalization-based therapy (MBT; Bateman & Fonagy, 2006). Randomized controlled trials (RCT; e.g., Bateman & Fonagy, 2009; Doering et al., 2010; Linehan et al., 2006) have shown that each of these interventions lead to significant reductions in BPD symptoms (e.g., suicidal behaviours and self-harm), compared with treatment as usual (for a review, see Stoffers et al., 2012).

Nonetheless, current psychotherapeutic interventions for BPD have important limitations, including limited treatment efficacy. For instance, compared with control interventions, a recent meta-analysis

[☆] This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

* Corresponding author. Department of Psychology, Ryerson University, 350 Victoria Street, Toronto, ON M5B 2K3, Canada.

E-mail address: rzeifman@psych.ryerson.ca (R.J. Zeifman).

(Cristea et al., 2017; $N = 2256$) found only small to moderate effect sizes for DBT (Hedges' $g = 0.34$) and psychodynamic approaches (i.e., TFP and MBT; Hedges' $g = 0.41$) on BPD outcomes (i.e., BPD symptoms, self-harm, and suicidal behaviors). Similarly, a recent longitudinal study showed that 10 years after treatment 45% of individuals continued to meet criteria for BPD and few showed improvement in social and occupational functioning (Alvarez-Tomás et al., 2017). Therefore, a need for improving the efficacy of psychotherapeutic interventions for BPD exists.

1.2. Pharmacological interventions for BPD

Compared with psychotherapeutic interventions, there has been less success developing efficacious pharmacological interventions for BPD. Currently, medical guidelines related to pharmacotherapy for BPD take two approaches: recommending against pharmacotherapy for BPD (e.g., National Institute of Health and Clinical Excellence, 2009) and recommending pharmacotherapy for specific BPD symptoms (e.g., APA, 2001). A small number of studies have explored adding pharmacological interventions to psychotherapeutic ones, with limited improvement in outcomes compared with psychotherapy alone (for a review, see Marin, Kapil-Prasad, Harris, & Goodman, 2018). Furthermore, meta-analyses and comprehensive reviews suggest that little evidence exists for the efficacy of current pharmacological interventions for BPD (Francois, Roth, & Klingman, 2015; Hancock-Johnson, Griffiths, & Picchioni, 2017; Lieb, Völlm, Rucker, Timmer, & Stoffers, 2010; Stoffers & Lieb, 2015), and no pharmacological interventions have health agency indication or approval for treatment of BPD (Starcevic & Janca, 2018).

Despite the limited evidence for pharmacological interventions for BPD, pharmacotherapy is frequently prescribed in the absence of a comorbid disorder (e.g., 82%; Paton, Crawford, Bhatti, Patel, & Barnes, 2015), suggesting that service providers are desperate to offer some form of treatment. Concerns with these practices include ethical issues, negative side effects, and high rates of medication abuse (Sansone & Wiederman, 2009). Given the limitations surrounding current interventions, a pressing need exists for exploring novel pharmacological interventions for BPD (Chanen, 2015).

2. Psychedelic-assisted psychotherapy

Given the need for novel pharmacological interventions for BPD, this review considers the potential of incorporating psychedelics within interventions for BPD. Broadly defined, psychedelics are pharmacological agents that cause changes in affect, cognition, and perception, as well as induce non-ordinary states of consciousness (dos Santos, Osorio, Crippa, & Hallak, 2016). Although there is some overlap between the phenomenological properties of psychedelics, many of them have different primary mechanisms of action. Serotonergic psychedelics (also referred to as "classic psychedelics"), such as psilocybin, dimethyltryptamine (DMT; which is contained in the ayahuasca brew), and lysergic acid diethylamide (LSD), share a common primary mechanism of action via agonism of serotonin 5-HT_{2A} receptors (Nichols, 2016). 3,4-Methylenedioxymethamphetamine (MDMA) is often defined as an entactogen (due to its modulation of feelings of interpersonal closeness and empathy; Bershad et al., 2019) and its mechanisms of action include increased levels of serotonin, dopamine, and norepinephrine (Sessa, Higbed, & Nutt, 2019). Ketamine is generally defined as a dissociative anaesthetic that acts as a glutamate *N*-methyl-D-aspartate (NMDA) receptor antagonist (Molero et al., 2018). Despite these important differences, given that this is the first review on psychedelics and BPD, we will explore the potential surrounding incorporating a broad class of psychedelics within interventions for BPD.

Preliminary research on the clinical utility of psychedelics has yielded promising results for treatment of a wide range of difficult to treat psychiatric disorders, a number of which are highly comorbid with

BPD (Zanarini, Frankenburg, Hennen, Reich, & Silk, 2004), including treatment-resistant major depressive disorder, posttraumatic stress disorder, substance use disorders, psychological distress associated with a life-threatening illness, and obsessive-compulsive disorder (for reviews, see Johnson, Hendricks, Barrett, & Griffiths, 2019; Sessa et al., 2019). A number of trials have also found that administration of psychedelics lead to long-lasting changes in personality (e.g., openness to experience; MacLean, Johnson, & Griffiths, 2011; Wagner et al., 2017), which may account for positive therapeutic outcomes (for a review, see Bouso, dos Santos, Alcázar-Córcoles, & Hallak, 2018). Furthermore, incorporating psychedelics within interventions for healthy individuals lead to long-lasting improvements in well-being (e.g., Griffiths et al., 2018; Nicholas et al., 2018). Despite important limitations surrounding research to date, including a limited number of trials, small sample sizes, and issues with maintaining blindness in placebo-controlled trials (due to the pronounced psychological effects of psychedelics; Barnby & Mehta, 2018), the preliminary results are promising and suggest that additional research on the clinical utility of psychedelics is warranted.

Given the wide range of psychiatric disorders for which psychedelics have shown promising results, Carhart-Harris, Erntzoe, Haijen, Kaalen, and Watts (2018) argue that psychedelics may target transdiagnostic mental health issues, which suggests that psychedelics may show promise for treatment of not yet researched psychiatric disorders. However, research to date has not yet explored the potential of incorporating psychedelics within interventions for BPD (and some research has actively excluded individuals with BPD; e.g., Danforth et al., 2018). Therefore, we review treatment targets in interventions for BPD, including (a) behavioral dysregulation, (b) emotion dysregulation, (c) disturbances in self-identity, and (d) social functioning, as well as the impact of psychedelics on these treatment targets. Furthermore, we describe research on purported overlapping mechanisms of change (i.e., how and why interventions lead to improvement; Kazdin, 2009) between interventions for BPD and psychedelics.

2.1. Behavioral dysregulation

2.1.1. Borderline personality disorder and behavioral dysregulation

BPD is characterized by a range of dysregulated behaviors including impulsivity, aggression, substance abuse, and suicidal behaviors (APA, 2013). Compared with healthy controls, individuals with BPD show higher levels of self-reported impulsivity (Sebastian, Jacob, Lieb, & Tüscher, 2013) and, among individuals with BPD, higher levels of impulsivity are associated with more frequent engagement in dysregulated behaviours (Terzi et al., 2017). Furthermore, among psychiatric inpatients, individuals with BPD are more likely to engage in serious violent and aggressive acts (Newhill, Eack, & Mulvey, 2009). BPD is also associated with high rates of comorbid lifetime alcohol (i.e., 48.1%) and drug (i.e., 45.9%) use disorders (Trull et al., 2016), as well as extremely high rates of non-suicidal self-injury (i.e., 65–80%; Clarkin, Widiger, Frances, Hurt, & Gilmore, 1983; Soloff, Lis, Kelly, Cornelius, & Ulrich, 1994), attempted suicide (e.g., 70.6%–83.6%; Soloff, Lynch, & Kelly, 2002; Wilson, Fertuck, Kwitel, Stanley, & Stanley, 2006), and completed suicide (i.e., 5.9%; Ternes et al., 2019). Thus, behavioral dysregulation is an important target in interventions for BPD.

Although evidence-based interventions for BPD lead to significant decreases in behavioral dysregulation, important limitations with treatment outcomes remain. For instance, comparing DBT with waitlist control and treatment as usual, a recent meta-analysis found only small to moderate effect sizes for decreases in self-harm and suicidal behaviors (Cohen's $d = -0.324$; DeCou, Comtois, & Landes, 2019) and non-significant decreases in suicidal ideation ($d = -0.229$). Moreover, following treatment of BPD, the majority of individuals (e.g., 58%; Heath, Laporte, Paris, Hamdullahpur, & Gill, 2018) show minimal improvement in substance abuse. Given these limitations, a need for developing interventions for BPD that better target behavioral

dysregulation exists.

2.1.2. Psychedelics and behavioral dysregulation

Research suggests that psychedelics may show potential for reducing behavioral dysregulation. Interviewed patients report increased behavioral control after psilocybin-assisted psychotherapy (Bogenschutz et al., 2018). Similarly, among healthy subjects, a single dose of LSD led to increases in self-reported positive behavioral changes 1 and 12 months later (Schmid & Liechti, 2018). Additionally, an observational study found significant decreases in self-reported impulsivity after 3–9 months of biweekly ritual ayahuasca use (Fernández et al., 2014). Psychedelics are also associated with decreases in aggression in both humans (for a review, see Tomlinson, Brown, & Hoaken, 2016) and non-human animals (e.g., Kostowski, Rewerski, & Piechocki, 1972). For instance, lifetime psychedelic use is associated with less likelihood of perpetrating a violent crime (Hendricks et al., 2017) and intimate partner violence (Thiessen, Walsh, Bird, & Lafrance, 2018; Walsh et al., 2016).

Research also suggests that psychedelics may decrease substance abuse. A meta-analysis found that treatment with LSD led to significant improvement among individuals with alcohol dependence (Krebs & Johansen, 2012). Small open-label studies have also found promising effects for treatment of tobacco (Johnson, Garcia-Romeu, Cosimano, & Griffiths, 2014) and alcohol dependence (Bogenschutz et al., 2015) with psilocybin-assisted psychotherapy. Additionally, lifetime use of psychedelics is associated with reduced risk of past-year opioid abuse (27%) and opioid dependence (40%; Pisano et al., 2017).

Psychedelics may lead to decreases in suicidal behaviors. A wide range of research has shown support for the rapid-acting anti-suicidal effects of ketamine (for a review, see Molero et al., 2018). Importantly, two of these studies included individuals with BPD (Grunebaum et al., 2017; Murrough et al., 2015) and BPD was not found to impact treatment outcomes (Grunebaum et al., 2017). Furthermore, among individuals with treatment-resistant depression, two doses of psilocybin with psychological support led to reductions in self-reported suicidal ideation 1 and 2 weeks after the intervention (Carhart-Harris, Bolstridge et al., 2018). Similarly, clinical trials have found that psilocybin-assisted psychotherapy and administration of ayahuasca lead to reductions in predictors of suicidal behaviors (for a review, see Johnson et al., 2019), such as depressive symptoms and hopelessness. Finally, lifetime use of psychedelics is associated with a reduced risk of becoming suicidal (Argento, Braschel, Walsh, Socias, & Shannon, 2018; Argento et al., 2017) and lower levels of past-year suicide ideation, planning, and attempts (Hendricks, Thorne, Clark, Coombs, & Johnson, 2015). However, these results may also be due to factors associated with psychedelic use (e.g., openness to experience), rather than use of psychedelics themselves. Overall, this research suggests that administration of psychedelics leads to decreases in a wide range of dysregulated behaviors associated with BPD.

3. Emotion dysregulation

3.1. Borderline personality disorder and emotion dysregulation

The core dysfunction of BPD is often identified as emotion dysregulation (Carpenter & Trull, 2013; Linehan, 1993). Emotion dysregulation is a multidimensional construct that consists of heightened negative affect, emotional sensitivity, and deficits in using emotion regulation strategies (Carpenter & Trull, 2013). Current neurobiological research suggests that emotion dysregulation among individuals with BPD stems from increased activity in limbic regions of the brain (e.g., amygdala, insula), which are associated with emotion detection and generation, coupled with decreased activity in prefrontal regions (e.g., prefrontal cortex), which are associated with emotion regulation (Schulze, Schmahl, & Niedtfeld, 2016). Therefore, emotion dysregulation is thought to occur due to deficits in “top-down” modulation of

emotional responses.

Given its centrality to BPD, emotion dysregulation is a primary target and purported mechanism of change in interventions for BPD (Rizvi & Thomas, 2017; Rudge, Feigenbaum, & Fonagy, 2017). Changes in self-reported emotion dysregulation, and related processes, mediate BPD treatment outcomes (Axelrod, Perepletchikova, Holtzman, & Sinha, 2011; Gratz, Bardeen, Levy, Dixon-Gordon, & Tull, 2015, 2012; McMain et al., 2013). Research also suggests that psychotherapeutic interventions for BPD alter the function, structure, and connectivity of regions of the brain essential to emotion dysregulation (for a review, see Herpertz, Schneider, Schmahl, & Bertsch, 2018). Overall, these findings suggest that emotion dysregulation is likely a key mechanism of change in interventions for BPD.

3.2. Psychedelics and Emotion Dysregulation

Empirical and theoretical research suggests that psychedelics impact emotional processes and emotion dysregulation (Vollenweider & Kometer, 2010). First, in a recent observational study with a community sample (Domínguez-Clavé et al., 2019), administration of ayahuasca led to significant decreases in components of emotion dysregulation (i.e., emotional non-acceptance, emotional interference, and lack of control) 24 h after administration. Within a subgroup of this sample, individuals high in BPD traits also showed decreases in components of emotion dysregulation (i.e. emotional interference and lack of control). Second, within community samples, lifetime psychedelic use is associated with lower levels of emotion dysregulation (Thiessen et al., 2018) and self-reported psychological distress (Hendricks et al., 2015). Third, research on psychedelic-assisted psychotherapy has shown long-lasting (e.g., 6–6.5 months) decreases in affective symptoms (for a review, see Johnson et al., 2019), as well as decreases in neuroticism (Erritzoe et al., 2018). Among individuals with a broad range of affective disorders, those that received high and low dose LSD showed greater improvement in affective symptoms than those who received conventional treatment alone (e.g., psychotropic medication, electroshock therapy, individual psychotherapy or group psychotherapy; McCabe, Savage, Kurland, & Unger, 1972). Furthermore, among five individuals with BPD and comorbid treatment-resistant depression who received ayahuasca, all five individuals showed decreases (i.e., 30–84% decrease from baseline) in clinician-assessed symptoms of depression 7 days later (Palhano-Fontes et al., 2019).

Administration of ketamine has also been shown to lead to rapid reductions in affective symptoms (for a review, see Molero et al., 2018). Interestingly, compared with individuals with low symptom burden at baseline (e.g., lower levels of suicidality), ketamine-assisted psychotherapy leads to greater decreases in affective symptoms among individuals higher in symptom burden (e.g., higher levels of suicidality; Dore et al., 2019). Finally, individuals with BPD and alcohol use disorder who received ketamine-assisted psychotherapy, showed improvements in characteristics related to emotion dysregulation, including measures related depressive symptoms and neuroticism (Krupitsky & Grinenko, 1996).

The impact of psychedelics on emotion dysregulation may be accounted for by the agonist action of psychedelics on 5-HT_{2A} receptors in the pre-frontal cortex, which lead to enhanced “top-down” control of circuits related to emotion dysregulation (for a review, see Vollenweider & Kometer, 2010). In line with this hypothesis, experimental research with humans indicates that psychedelics alter neural activity in regions associated with emotion processing and emotion dysregulation (e.g., Bedi, Phan, Angstadt, & De Wit, 2009; Kraehenmann et al., 2016; Sanches et al., 2016). Furthermore, administration of psychedelics has been shown to promote structural and functional plasticity, and related effects (i.e., increased gene transcription), in the pre-frontal cortex of rats (Ly et al., 2018; Nichols & Sanders-Bush, 2002). Therefore, preliminary research suggests that psychedelics may target emotion dysregulation.

4. Mindfulness

4.1. Borderline personality disorder and mindfulness

One important emotion regulation strategy relevant to BPD is mindfulness, and related capacities, including decentering (i.e., the ability to observe one's thoughts, emotions, and sensation from a detached perspective; Kerr, Josyula, & Littenberg, 2011) and acceptance (i.e., tolerating and approaching, rather than avoiding, one's experiences; Orzech, Shapiro, Brown, & McKay, 2009). Compared with individuals with other personality disorders (Zanarini, Temes, Frankenburg, Reich, & Fitzmaurice, 2019), as well as community (e.g., Tortella-Feliu et al., 2018) and college (Baer, Smith, & Allen, 2004) comparison groups, individuals with BPD show lower levels of mindfulness and acceptance. Moreover, among individuals with BPD, mindfulness is positively associated with emotional well-being and negatively associated with healthcare utilization (O'Toole, Diddy, & Kent, 2012). Furthermore, within a psychiatric sample, deficits in mindfulness account for the relationship between BPD features and behavioral dysregulation (Wupperman, Fickling, Klemanski, Berking, & Whitman, 2013). Thus, deficits in mindfulness play an important role in BPD.

Mindfulness is purported to be a key target and mechanism of change in interventions for BPD (Bliss & McCardle, 2014; Linehan, 1993; Rudge et al., 2017). Time spent engaging in mindfulness practice (Feliu-Soler et al., 2014; Soler et al., 2012), use of mindfulness skills (Stepp, Epler, Jahng, & Trull, 2008), as well as increases in mindfulness (Mitchell, Roberts, Bartsch, & Sullivan, 2019; Zeifman, Boritz, Barnhart, Labrish, & McMain, 2019), acceptance (Krantz, McMain, & Kuo, 2018; Perroud, Nicastro, Jermann, & Huguelet, 2012; Zanarini et al., 2019), and decentering (Elices et al., 2016) are associated with positive BPD treatment outcomes. Finally, increases in mindfulness throughout interventions for BPD are associated with less likelihood of treatment drop out (Stratton, Alvarez, Labrish, Barnhart, & McMain, 2018). These results suggest that mindfulness may be an especially important mechanism of change in interventions for BPD. Therefore, pharmacological interventions that increase mindfulness may strengthen BPD treatment outcomes.

4.2. Psychedelics and Mindfulness

Mindfulness has been proposed as a mechanism of change in psychedelic-assisted psychotherapy. Qualitative research has highlighted mindfulness (Bogenschutz et al., 2018) and acceptance (Wagner, Mithoefer, Mithoefer, & Monson, 2019; Watts, Day, Krzanowski, Nutt, & Carhart-Harris, 2017) as important processes of change associated with psilocybin and MDMA-assisted psychotherapy. Furthermore, experimental studies show that administration of ayahuasca leads to increases in mindfulness-related capacities, including acceptance and decentering (Domínguez-Clavé et al., 2019; Sampedro et al., 2017; Soler et al., 2018, 2016; Thomas, Lucas, Capler, Tupper, & Martin, 2013). In one ayahuasca study, participants continued to show significant improvement in acceptance at 2-month follow-up (Sampedro et al., 2017). Similarly, Soler et al. (2018) found that four sessions of ayahuasca result in increases in acceptance that are comparable with 8-weeks of mindfulness-based stress reduction (Kabat-Zinn, 1990). Finally, lifetime use of ayahuasca is associated with higher levels of decentering (Franquesa et al., 2018). Overall, these results suggest that administration of psychedelics may lead to increases in mindfulness.

5. Self-identity

5.1. Borderline personality disorder and self-identity

Individuals with BPD experience disturbances in their self-identity (APA, 2013), which have been conceptualized in two primary ways

(Gad et al., 2019): (a) as an inconsistent and non-integrated sense of self (Kernberg, 1975) and (b) as a generally elevated negative view toward the self with only brief periods of positive views (Zanarini et al., 2007). Research has provided support for both of these models. For instance, compared with healthy controls, individuals with BPD show less consistency when listing their personality traits (Beeney, Hallquist, Ellison, & Levy, 2016), as well as less coherent and more disoriented life narratives (Jørgensen et al., 2012). Furthermore, compared with healthy controls and clinical populations, individuals with BPD show heightened negative self-evaluation and self-criticism, as well as less self-compassion (e.g., Costa, Marôco, Pinto-Gouveia, Ferreira, & Castilho, 2016; Gad et al., 2019; Kopala-Sibley, Zuroff, Russell, Moskowitz, & Paris, 2012; Winter, Bohus, & Lis, 2017). Moreover, self-criticism and self-compassion are associated with maladaptive behaviors associated with BPD, including self-injury and suicidal behaviors (Gregory, Glazer, & Berenson, 2017; Hasking, Boyes, Finlay-Jones, McEvoy, & Rees, 2019).

Qualitative analysis of interviews conducted with BPD treatment providers and users identify self-compassion as an important theme in the process of recovery (Katsakou & Pistrang, 2017; Katsakou, Pistrang, Barnicot, White, & Priebe, 2017). Research has also suggested that incorporating self-compassion-based exercises improves outcomes in interventions for BPD (Feliu-Soler et al., 2017). Furthermore, compared with BPD non-responders, treatment responders show lower levels of baseline negative self-views and greater decreases in negative self-views (Gad et al., 2019), as well as steeper increases in self-acceptance (Zanarini et al., 2019). Finally, a cross-sectional study showed that, after treatment for BPD, personal recovery was negatively associated with self-criticism and positively associated with self-compassion (Donald, Lawrence, Broadbear, & Rao, 2019). Accordingly, disturbances in self-identity, including instability of views toward the self, as well as increasing positive views toward the self and increasing self-compassion are important, and especially challenging, treatment targets in interventions for BPD (Krawitz, 2012).

5.2. Psychedelics and self-identity

Research suggests that psychedelics have an acute, and potentially long-lasting, impact on an individual's sense of identity. Qualitative research indicates that self-compassion and self-acceptance are important themes that emerge from interviews with individuals that received psilocybin, LSD, and MDMA-assisted psychotherapy, or used ayahuasca for therapeutic purposes (Barone, Beck, Mitsunaga-Whitten, & Perl, 2019; Bogenschutz et al., 2018; Gasser, Kirchner, & Passie, 2015; Lafrance et al., 2017; Malone et al., 2018). Cross-sectional and experimental research similarly suggests that psychedelics lead to increased self-compassion and positive views of the self. Among healthy individuals, controlled studies with psilocybin (Griffiths, Richards, McCann, & Jesse, 2006, 2008, 2011, 2018), and an open-label study with LSD (Schmid & Liechti, 2018), have shown long-term increases in positive views toward the self. Relatedly, compared with individuals without a history of ayahuasca use, individuals with a history of ayahuasca use show more positive views of the self (Franquesa et al., 2018). An open-label study with healthy volunteers showed that a single dose of ayahuasca leads to significant increases in self-compassion (Sampedro et al., 2017). Similarly, among recreational MDMA users, use of MDMA is associated with subsequent increases in self-compassion and decreases in self-criticism (Kamboj et al., 2015, 2017). Among individuals with BPD and substance use, ketamine-assisted psychotherapy was found to lead improvement in personality characteristics associated with self-criticism (Krupitsky & Grinenko, 1996). These findings suggest that administration of psychedelics may lead to long-lasting changes in self-identity in general and increases in self-compassion in particular.

To date, research has not yet explored the impact of psychedelics on stability of views toward the self (i.e., BPD-related identity disturbance

as defined by Kernberg, 1975). However, given research suggesting that decreased emotion dysregulation (Gratz et al., 2015) and use of mindfulness skills (Stepp et al., 2008) lead to decreases in disturbance in self-identity within interventions for BPD, the impact of psychedelics on emotion dysregulation (see above, ‘**Psychedelics and Emotion Dysregulation**’) and mindfulness (see above, ‘**Psychedelics and Mindfulness**’) may, in turn, lead to decreases in disturbances in self-identity. Additional research will be necessary to explore this possibility.

6. Social functioning

6.1. Borderline personality disorder and social functioning

BPD is characterized by deficits in social functioning, including unstable and intense interpersonal relationships, and frantic efforts to avoid abandonment (APA, 2013). Individuals with BPD are especially likely to experience heightened levels of social pain (i.e., emotional distress due to social exclusion, rejection, or loss; Renneberg et al., 2012; Staebler, Renneberg et al., 2011) and rejection sensitivity (Domsalla et al., 2014; Staebler, Helbing, Rosenbach, & Renneberg, 2011b). Furthermore, individuals with BPD may show additional deficits in aspects of social functioning, including trust, emotion recognition, and empathy (for a review, see Lazarus, Cheavens, Festa, & Rosenthal, 2014).

Deficits in social functioning are among the most difficult targets in BPD interventions, with little evidence for increases in social functioning following treatment (Alvarez-Tomás et al., 2017; Gunderson et al., 2011). One of the means through which deficits in social functioning can manifest itself in interventions for BPD is through interference with the therapeutic alliance (i.e., the relationship between the patient and therapist). Maintenance of therapeutic alliance can be especially difficult (Bender, 2005), and is an important predictor of drop-out and treatment outcomes in interventions for BPD (Barnicot et al., 2012; Barnicot, Katsakou, Marougka, & Priebe, 2011; Bedics, Atkins, Harned, & Linehan, 2015; Wnuk et al., 2013). Therefore, improving social functioning, as well as enhancement of the therapeutic alliance, are important targets in interventions for BPD.

6.2. Psychedelics and social functioning

Research suggests that psychedelics may have a positive impact on social functioning. Among individuals with BPD and substance use, ketamine-assisted psychotherapy led to improvements in social functioning and interpersonal sensitivity (Krupitsky & Grinenko, 1996). Furthermore, among individuals that received psilocybin with psychological support, thematic analysis identified connectedness to others as an important process of change (Watts et al., 2017). Similarly, after psilocybin and MDMA-assisted psychotherapy, individuals reported experiencing improvements in their relationships (Barone, Beck, Mitsunaga-Whitten, & Perl, 2019; Belser et al., 2017). Furthermore, in an RCT, healthy individuals receiving standard support for spiritual practice and two high doses of psilocybin (vs. two low doses) showed increases in closeness to others (Griffiths et al., 2018). Moreover, in healthy participants, a single dose of LSD leads to increases in positive social effects 1 and 12 months later (Schmid & Liechti, 2018). Finally, within a community sample, use of psychedelics leads to increased social connectedness 2 weeks later (Carhart-Harris, Erritzoe et al., 2018). In line with these findings, Carhart-Harris, Erritzoe et al. (2018) argue that connectedness may be a core transdiagnostic issue that is targeted by psychedelics.

Psychedelics may also target deficits in social functioning that are associated with BPD, including social pain, rejection sensitivity, trust, empathy, and emotion recognition. For instance, among individuals with treatment-resistant depression, psilocybin with psychological support improved emotional face recognition two weeks after

administration, such that they were no longer different from healthy controls (Stroud et al., 2018). Similarly, double-blind randomized controlled experiments with healthy controls have shown that administration of psilocybin, LSD, and MDMA acutely reduce feelings of social pain, social exclusion, and recognition of negative emotions, and increase emotional empathy, trust, and closeness to others (e.g., Dolder, Schmid, Müller, Borgwardt, & Liechti, 2016; Frye, Wardle, Norman, & de Wit, 2014; Gabay, Kempton, Gilleen, & Mehta, 2019; Hysek et al., 2013; Komater et al., 2012; Pokorny, Preller, Komater, Dziobek, & Vollenweider, 2017; Preller et al., 2015, 2016). Based on these results, it has been suggested that psychedelics may be used to target psychiatric disorders characterized by social deficits, as well as enhance therapeutic alliance, which are important targets within interventions for BPD.

7. Risk, safety, and clinical considerations

Research suggests that psychedelics are safe when administered within controlled environments after careful screening. Serotonergic psychedelics show low levels of physiological toxicity, are not associated with bodily harm, neuropsychological deficits, or mental health problems and show low abuse and dependence potential (Hamill, Hallak, Dursun, & Baker, 2019; Johnson, Griffiths, Hendricks, & Henningfield, 2018; Johnson, Richards, & Griffiths, 2008), which is important given high rates of substance abuse among individuals with BPD (Trull et al., 2016). Mental health concerns very rarely occur (i.e., < 0.2% in vulnerable populations) following administration of serotonergic psychedelics in controlled settings (Cohen & Ditman, 1963). Among over 2000 patients that have been administered psilocybin, LSD, and ayahuasca in controlled research settings, there have been no serious adverse events reported (dos Santos et al., 2016; Johnson et al., 2008). Although MDMA is associated with greater abuse liability than serotonergic psychedelics, MDMA has been identified as being less dangerous than other frequently prescribed pharmacological agents (e.g., benzodiazepines; Nutt, King, & Phillips, 2010) and use of MDMA in a non-medical setting is rare after its administration in clinical contexts (Mithoefer et al., 2013). MDMA has been administered more than 1600 times within controlled settings and only one serious adverse event (i.e., a brief increase in pre-existing ventricular ectopy) and no deaths have been reported (Multidisciplinary Association for Psychedelics Studies, 2017).

The safety profile of ketamine is less clear, with potential side effects including dependence, craving, cognitive, urological, and hepatic concerns (for a review, see Strong & Kabbaj, 2018). Nonetheless, within clinical contexts, administration of ketamine has not been associated with persistent medical or mental health issues (e.g., substance abuse; Dore et al., 2019) and has been approved for a range of clinical indications, including treatment-resistant depression. As with all pharmacological interventions, important considerations exist related to the potential for negative interactions between psychedelics and other medications (e.g., serotonin reuptake inhibitors), as well as physical health issues (e.g., high blood pressure; for guidelines, see, Johnson et al., 2008; Kim & Mierzwinski-Urban, 2017; Sessa et al., 2019).

The primary issues related to administration of psychedelics are the potential for acute physiological discomfort (e.g., dizziness, weakness, tremors, nausea, vomiting, paraesthesia, blurred vision, and increased tendon reflexes) and challenging psychological experiences (e.g., anxiety, fear, and confusion; Hamill et al., 2019; Nichols, 2004). These challenging psychological experiences are generally considered an important component of the psychotherapeutic process (e.g., Carbonaro et al., 2016) and can generally be managed within clinical settings through preparation, session monitoring, and integration of challenging experiences.

Limited data regarding the safety and tolerability of psychedelics for individuals with BPD exists. A search of the literature identified nine studies that administered a psychedelic to individuals with BPD,

including a 3–9 month study of biweekly ritual ayahuasca administration that included one individual with BPD (Fernández et al., 2014), a single dose of ayahuasca to five individuals with BPD (Palhano-Fontes et al., 2019), and one to six administrations of ketamine to 63 individuals with BPD (Aust et al., 2019; Cole et al., 2018; Feifel, Malcolm, Boggie, & Lee, 2017; Grunebaum et al., 2017; Krupitsky & Grinenko, 1996; Lauritsen, Mazuera, Lipton, & Ashina, 2016; Murrrough et al., 2015). Importantly, no serious adverse events were reported related to administration of psychedelics in any of these studies. Interestingly, Peter Gasser who has been conducting therapy with psychedelics in Switzerland for several decades, notes that “From 1988 to 1993, a significant number of patients with narcissistic personality disorders sought therapy with psychedelic drugs. The borderline personality disorder was also diagnosed rather often, as were depressed mood disorders and adjustment disorders. We can presume that the treatment is well suited for these disorders (Gasser, 1994, p. 7).”

It is important to note that Carhart-Harris, Bolstridge et al. (2018) report that they plan to exclude individuals with BPD from future trials with psilocybin for treatment of depression (see [clinicaltrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT03429075); NCT 03429075). Similarly, past MDMA-assisted psychotherapy trials have excluded individuals with BPD (e.g., Danforth et al., 2018; Mithoefer, Wagner, Mithoefer, Jerome, & Doblin, 2011). Furthermore, other trials have excluded “psychiatric conditions judged to be incompatible with establishment of rapport or safe exposure to psilocybin (e.g., Griffiths et al., 2016, Supplementary material).” Additionally, some trials have excluded individuals with current serious suicide risk (e.g., Palhano-Fontes et al., 2019) or substance abuse (e.g., Ross et al., 2016), which may have indirectly resulted in exclusion of certain individuals with BPD.

Given the complexity surrounding BPD, the short-term nature of these trials, and that other psychiatric disorders were the primary interest, these decisions are understandable. Nonetheless, it is important to note that clinical trials with psychedelics have reported including individuals with current serious suicidal ideation and past suicidal behaviors (e.g., O’alora et al., 2018). Furthermore, current safety guidelines for conducting clinical research with serotonergic psychedelics do not specifically mention exclusion of individuals with BPD (although they do suggest excluding individuals with risk of psychotic disorders, as well as bipolar I or II disorder; Johnson et al., 2008).

The primary safety concerns related to administration of psychedelics are during their acute effects. Given the association between BPD and suicidal behaviors, it is especially important that any potential safety risks be minimized. Therefore, in line with common practice in psychiatric trials (e.g., Linehan, McDavid, Brown, Sayrs, & Gallop, 2008), we recommend that initial studies on the safety and tolerability of psychedelics for individuals with BPD exclude individuals with recent (i.e., in the past 8 weeks) serious self-harm or suicidal behaviors. Furthermore, in line with treatment guidelines for psychedelic-assisted psychotherapy (Johnson et al., 2008), it is recommended that administration of psychedelics occur (a) within controlled settings in which safety can be ensured, (b) in the presence of a treatment provider that can provide support, guidance and, although generally unnecessary, medication (e.g., benzodiazepines) if needed, (c) in a warm and comfortable environment, accompanied by the opportunity to listen to a selectively chosen music playlist (for the important role of music in psychedelic-assisted psychotherapy, see Kaelen et al., 2018), and (d) that preparation and integration sessions prior to and following administration of psychedelics be provided. Furthermore, during the preparation and integration sessions, we suggest that patients be encouraged to take time to integrate their experiences before making any major life decisions or changes (e.g., quitting their job, ending a marriage). Given the importance of the therapeutic alliance within psychedelic-assisted psychotherapy, as well as the potential difficulty surrounding establishing a strong therapeutic alliance with individuals with BPD (Bender, 2005), it is important psychedelic sessions occur only after patients report trusting their therapist. Additionally, with

serotonergic psychedelics and ketamine, it may be helpful to begin with small doses and increase the dosage as the therapeutic alliance and commitment to change develop.

Given the complexity associated with BPD, we recommend that psychedelic-assisted psychotherapy be incorporated within an already established evidence-based intervention for BPD (e.g., DBT, TFP, or MBT). Relatedly, a recent study incorporating MDMA-assisted psychotherapy within an adapted version of DBT showed promising results for treatment of autistic adults with social anxiety disorder (Danforth et al., 2018). There is substantial overlap between BPD treatment targets and psychedelics. The principles of DBT and psychedelic-assisted psychotherapy show substantial overlap, including emphases on emotion dysregulation and enhancing mindfulness (for a discussion, see Walsh & Thiessen, 2018). There is also evidence that combining psychedelics with meditation, an essential component of DBT (Linehan, 1993), has positive and long-lasting synergistic effects (e.g., Smigielski, Scheidegger, Kometer, & Vollenweider, 2019). Furthermore, given the emphasis on behavioural change within DBT (Linehan, 1993) and suggestions that psychedelics induce a heightened state of plasticity (i.e., enhanced capacity to change; Carhart-Harris & Friston, 2019; Carhart-Harris & Nutt, 2017), integrating psychedelics and DBT may allow for optimal levels of behavioural change. Given the conceptual overlap between mentalization (i.e., making sense of one’s social world by understanding one’s own and others mental states) and mindfulness (Choi-Kain & Gunderson, 2008), integrating psychedelics with MBT may similarly help to optimize improved mentalization capacities. Finally, it has been suggested that psychedelics facilitate experiences of transference (Goldsmith, 2007) and enhance therapeutic alliance (e.g., Dolder et al., 2016; Mithoefer et al., 2011), which are essential components of MBT and TFP. Moreover, the altered sense of self (for a discussion, see Nour & Carhart-Harris, 2017) and relation to others (Bershad et al., 2019; Carhart-Harris, Erritzoe et al., 2018) induced by psychedelics and the emphasis on targeting the patient’s representation of the self and others within TFP (Clarkin, Cain, & Lenzenweger, 2018), may benefit from one another. Accordingly, compared with either treatment alone, utilizing psychedelics alongside evidence-based interventions and their techniques, such as DBT (e.g., mindfulness, radical acceptance), MBT (e.g., ‘validation of the transference,’ ‘mentalizing the transference’), and TFP (e.g., ‘working with transference,’ clarification, confrontation), may have positive synergistic effects for the treatment of BPD (see Fig. 1). Therefore, we argue that integrating evidence-based interventions for BPD with psychedelics will be essential for ensuring treatment safety and optimizing therapeutic outcomes.

8. Limitations and future research

The research described in this review has a number of important limitations. Much of the research described is cross-sectional, includes small sample sizes, fails to explore the long-term impact of psychedelics, and are based on open-label trials. Furthermore, maintaining blindness in placebo-controlled psychedelic research remains challenging (Barnby & Mehta, 2018). Most importantly, the research described in this review was not conducted with individuals with BPD. Therefore, in order to determine the safety, tolerability, and clinical utility of incorporating psychedelics within interventions for BPD, future research should aim to (a) establish the safety and tolerability of administering psychedelics to individuals with BPD, (b) explore the impact of psychedelics on behavioral dysregulation, emotion dysregulation, disturbances in self-identity, and social functioning among individuals with BPD, and (c) determine whether incorporating psychedelics within interventions for BPD improves treatment outcomes. Until such studies have been conducted, caution surrounding administration of psychedelics to individuals with BPD is strongly recommended.

One important barrier to conducting research with psychedelics are federal regulations (Nutt, King, & Nichols, 2013), which currently classify psychedelics, such as psilocybin, LSD, DMT, and MDMA as




Fig. 1. Psychedelic-assisted psychotherapy targets, components of BPD interventions, and BPD treatment targets.

Schedule 1 drugs and ketamine as a Schedule 3 drug. Nonetheless, approval for conducting clinical research with psychedelics has been granted for a range of psychiatric disorders. Moreover, MDMA, psilocybin, and esketamine have received “Breakthrough Therapy Designation” by the US Food and Drug Administration (FDA) for posttraumatic stress disorder, treatment-resistant depression, and major depressive disorder with imminent risk of suicide, respectively. Furthermore, esketamine being approved by the FDA for treatment-resistant depression may allow for additional research on its impact on individuals with comorbid BPD. Interestingly, an ongoing phase 2 clinical trial is exploring ketamine (vs. Midazolam) as an intervention for BPD (single dose, with no indication of including concurrent psychotherapy; clinicaltrials.gov, NCT 03395314). Therefore, although challenging, conducting research with psychedelics appears possible and may ultimately contribute to improving interventions for BPD.

9. Conclusion

In summary, BPD is a severe psychiatric disorder characterized by

behavioral dysregulation, emotion dysregulation, disturbances in self-identity, and deficits in social functioning. Despite the existence of evidence-based interventions for BPD, important limitations surrounding current interventions remain, including limited psychotherapeutic treatment efficacy (i.e., small-moderate effect sizes) and limited evidence for the efficacy of pharmacological interventions. Therefore, there exists a need for improving current interventions for BPD. Psychedelics, including psilocybin, ayahuasca, LSD, MDMA, and ketamine, show promise for treatment of a wide range of psychiatric disorders. Furthermore, research suggests that psychedelics may positively impact BPD treatment targets (i.e., behavioral dysregulation, emotion dysregulation, disturbances in self-identity, and social functioning) and target overlapping mechanisms of change in interventions for BPD (i.e., emotion dysregulation, mindfulness, and self-compassion). However, research has not yet directly explored the impact of psychedelics on BPD. Given safety considerations surrounding administration of psychedelics to individuals with BPD, we recommend strict guidelines, including administration of psychedelics within controlled and safe settings. Overall, this review suggests that conducting research on

incorporating psychedelics within therapeutic interventions for BPD (e.g., DBT) is warranted. Ultimately, research will be necessary to determine the clinical utility of incorporating psychedelics within interventions for BPD.

References

- Alvarez-Tomás, I., Soler, J., Bados, A., Martín-Blanco, A., Elices, M., Carmona, C., & Pascual, J. C. (2017). Long-term course of borderline personality disorder: A prospective 10-year follow-up study. *Journal of Personality Disorders*, 31, 590–605.
- American Psychiatric Association. (2001). *Practice guidelines for the treatment of patients with borderline personality disorder*. Washington, DC: American Psychiatric Association.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Association Publishing.
- Argento, E., Bräschel, M., Walsh, Z., Socias, M. E., & Shannon, K. (2018). The moderating effect of psychedelics on the prospective relationship between prescription opioid use and suicide risk among marginalized women. *Journal of Psychopharmacology*, 32, 1385–1391.
- Argento, E., Strathdee, S. A., Tupper, K., Bräschel, M., Wood, E., & Shannon, K. (2017). Does psychedelic drug use reduce risk of suicidality? Evidence from a longitudinal community-based cohort of marginalized women in a Canadian setting. *British Medical Journal Open*, 7, e016025.
- Aust, S., Gärtner, M., Basso, L., Otte, C., Wingenfeld, K., Chae, W. R., ... Grimm, S. (2019). Anxiety during ketamine infusions is associated with negative treatment responses in major depressive disorder. *European Neuropsychopharmacology*, 29, 529–538.
- Axelrod, S. R., Perepletchikova, F., Holtzman, K., & Sinha, R. (2011). Emotion regulation and substance use frequency in women with substance dependence and borderline personality disorder receiving dialectical behavior therapy. *The American Journal of Drug and Alcohol Abuse*, 37, 37–42.
- Baer, R. A., Smith, G. T., & Allen, K. B. (2004). Assessment of mindfulness by self-report: The Kentucky inventory of mindfulness skills. *Assessment*, 11, 191–206.
- Barnby, J. M., & Mehta, M. A. (2018). Psilocybin and mental health—don't lose control. *Frontiers in Psychiatry*, 9. <https://doi.org/10.3389/fpsy.2018.00293>.
- Barnicot, K., Katsakou, C., Bhatti, N., Savill, M., Fearn, N., & Priebe, S. (2012). Factors predicting the outcome of psychotherapy for borderline personality disorder: A systematic review. *Clinical Psychology Review*, 32, 400–412.
- Barnicot, K., Katsakou, C., Marougka, S., & Priebe, S. (2011). Treatment completion in psychotherapy for borderline personality disorder—a systematic review and meta-analysis. *Acta Psychiatrica Scandinavica*, 123, 327–338.
- Bateman, A. W., & Fonagy, P. (2006). *Mentalization based treatment for borderline personality disorder: A practical guide*. Oxford, UK: Oxford University Press.
- Barone, W., Beck, J., Mitsunaga-Whitten, M., & Perl, P. (2019). Perceived benefits of MDMA-assisted psychotherapy beyond symptom reduction: Qualitative follow-up study of a clinical trial for individuals with treatment-resistant PTSD. *Journal of Psychoactive Drugs*, 51, 199–208. <https://doi.org/10.1080/02791072.2019.1580805>.
- Bateman, A., & Fonagy, P. (2009). Randomized controlled trial of outpatient mentalization-based treatment versus structured clinical management for borderline personality disorder. *American Journal of Psychiatry*, 166, 1355–1364.
- Bedics, J. D., Atkins, D. C., Harned, M. S., & Linehan, M. M. (2015). The therapeutic alliance as a predictor of outcome in dialectical behavior therapy versus non-behavioral psychotherapy by experts for borderline personality disorder. *Psychotherapy*, 52, 67–77.
- Bedi, G., Phan, K. L., Angstadt, M., & De Wit, H. (2009). Effects of MDMA on sociability and neural response to social threat and social reward. *Psychopharmacology*, 207, 73–83.
- Beeney, J. E., Hallquist, M. N., Ellison, W. D., & Levy, K. N. (2016). Self-other disturbance in borderline personality disorder: Neural, self-report and performance-based evidence. *Journal of Personality Disorders*, 7, 28–39.
- Belser, A. B., Agin-Lieb, G., Swift, T. C., Terrana, S., Devenot, N., Friedman, H. L., ... Ross, S. (2017). Patient experiences of psilocybin-assisted psychotherapy: An interpretative phenomenological analysis. *Journal of Humanistic Psychology*, 57, 354–388.
- Bender, D. S. (2005). The therapeutic alliance in the treatment of personality disorders. *Journal of Psychiatric Practice*, 11, 73–87.
- Bershad, A. K., Mayo, L. M., Van Hedger, K., McGlone, F., Walker, S. C., & de Wit, H. (2019). Effects of MDMA on attention to positive social cues and pleasantness of affective touch. *Neuropsychopharmacology*, 44, 1698–1705.
- Bliss, S., & McCardle, M. (2014). An exploration of common elements in dialectical behavior therapy, mentalization based treatment and transference focused psychotherapy in the treatment of borderline personality disorder. *Clinical Social Work Journal*, 42, 61–69.
- Bogenschutz, M. P., Forchimes, A. A., Pommy, J. A., Wilcox, C. E., Barbosa, P. C. R., & Strassman, R. J. (2015). Psilocybin-assisted treatment for alcohol dependence: A proof-of-concept study. *Journal of Psychopharmacology*, 29, 289–299.
- Bogenschutz, M. P., Podrebarac, S. K., Duane, J. H., Amegadzie, S. S., Malone, T. C., Owens, L. T., ... Mennenga, S. E. (2018). Clinical interpretations of patient experience in a trial of psilocybin-assisted psychotherapy for alcohol use disorder. *Frontiers in Pharmacology*, 9, 100. <https://doi.org/10.3389/fphar.2018.00100>.
- Bouso, J. C., dos Santos, R. G., Alcázar-Córcoles, M.Á., & Hallak, J. E. (2018). Serotonergic psychedelics and personality: A systematic review of contemporary research. *Neuroscience & Biobehavioral Reviews*, 87, 118–132.
- Cailhol, L., Thalamas, C., Garrido, C., Birmes, P., & Lapeyre-Mestre, M. (2015). Mental health service utilization among borderline personality disorder patients inpatient. *L'Encephale*, 41, 115–122.
- Carbonaro, T. M., Bradstreet, M. P., Barrett, F. S., MacLean, K. A., Jesse, R., Johnson, M. W., & Griffiths, R. R. (2016). Survey study of challenging experiences after ingesting psilocybin mushrooms: Acute and enduring positive and negative consequences. *Journal of Psychopharmacology*, 30, 1268–1278.
- Carhart-Harris, R. L., Bolstridge, M., Day, C. M. J., Rucker, J., Watts, R., Erritzoe, D. E., ... Rickard, J. A. (2018b). Psilocybin with psychological support for treatment-resistant depression: Six-month follow-up. *Psychopharmacology*, 235, 399–408.
- Carhart-Harris, R. L., Erritzoe, D., Haijen, E., Kaelen, M., & Watts, R. (2018a). Psychedelics and connectedness. *Psychopharmacology*, 235, 547–550.
- Carhart-Harris, R. L., & Friston, K. J. (2019). REBUS and the anarchic brain: Toward a unified model of the brain action of psychedelics. *Pharmacological Reviews*, 71, 316–344.
- Carhart-Harris, R. L., & Nutt, D. J. (2017). Serotonin and brain function: A tale of two receptors. *Journal of Psychopharmacology*, 31, 1091–1120.
- Carpenter, R. W., & Trull, T. J. (2013). Components of emotion dysregulation in borderline personality disorder: A review. *Current Psychiatry Reports*, 15, 1–8.
- Chanen, A. M. (2015). Borderline personality disorder in young people: Are we there yet? *Journal of Clinical Psychology*, 71, 778–791.
- Choi-Kain, L. W., & Gunderson, J. G. (2008). Mentalization: Ontogeny, assessment, and application in the treatment of borderline personality disorder. *American Journal of Psychiatry*, 165, 1127–1135.
- Clarkin, J. F., Cain, N. M., & Lenzenweger, M. F. (2018). Advances in transference-focused psychotherapy derived from the study of borderline personality disorder: Clinical insights with a focus on mechanism. *Current Opinion in Psychology*, 21, 80–85.
- Clarkin, J. F., Widiger, T. A., Frances, A., Hurt, S. W., & Gilmore, M. (1983). Prototypic typology and the borderline personality disorder. *Journal of Abnormal Psychology*, 92, 263–275.
- Clarkin, J. P., Yeomans, F. E., & Kernberg, O. F. (1999). *Transference-focused psychotherapy for borderline personality disorder patients*. New York, NY: Wiley.
- Cohen, S., & Ditman, K. S. (1963). Prolonged adverse reactions to lysergic acid diethylamide. *Archives of General Psychiatry*, 8, 475–480.
- Cole, J. B., Klein, L. R., Nystrom, P. C., Moore, J. C., Driver, B. E., Fryza, B. J., ... Ho, J. D. (2018). A prospective study of ketamine as primary therapy for prehospital profound agitation. *The American Journal of Emergency Medicine*, 36, 789–796.
- Costa, J., Marôco, J., Pinto-Gouveia, J., Ferreira, C., & Castilho, P. (2016). Validation of the psychometric properties of the Self-Compassion Scale. Testing the factorial validity and factorial invariance of the measure among borderline personality disorder, anxiety disorder, eating disorder and general populations. *Clinical Psychology and Psychotherapy*, 23, 460–468.
- Cristea, I. A., Gentili, C., Cotet, C. D., Palomba, D., Barbui, C., & Cuijpers, P. (2017). Efficacy of psychotherapies for borderline personality disorder: A systematic review and meta-analysis. *JAMA Psychiatry*, 74, 319–328.
- Danforth, A. L., Grob, C. S., Struble, C., Feduccia, A. A., Walker, N., Jerome, L., & Emerson, A. (2018). Reduction in social anxiety after MDMA-assisted psychotherapy with autistic adults: A randomized, double-blind, placebo-controlled pilot study. *Psychopharmacology*, 235, 3137–3148.
- DeCou, C. R., Comtois, K. A., & Landes, S. J. (2019). Dialectical behavior therapy is effective for the treatment of suicidal behavior: A meta-analysis. *Behavior Therapy*, 50, 60–72.
- Doering, S., Hörz, S., Rentrop, M., Fischer-Kern, M., Schuster, P., Benecke, C., ... Buchheim, P. (2010). Transference-focused psychotherapy v. treatment by community psychotherapists for borderline personality disorder: Randomised controlled trial. *The British Journal of Psychiatry*, 196, 389–395.
- Dolder, P. C., Schmid, Y., Müller, F., Borgwardt, S., & Liechti, M. E. (2016). LSD acutely impairs fear recognition and enhances emotional empathy and sociality. *Neuropsychopharmacology*, 41, 2638–2646.
- Dominguez-Clavé, E., Soler, J., Pascual, J. C., Elices, M., Franquesa, A., Valle, M., ... Riba, J. (2019). Ayahuasca improves emotion dysregulation in a community sample and in individuals with borderline-like traits. *Psychopharmacology*, 236, 573–580.
- Domsalla, M., Koppe, G., Niedteld, I., Vollstädt-Klein, S., Schmah, C., Bohus, M., & Lis, S. (2014). Cerebral processing of social rejection in patients with borderline personality disorder. *Social Cognitive and Affective Neuroscience*, 9, 1789–1797.
- Donald, F., Lawrence, K. A., Broadbear, J. H., & Rao, S. (2019). An exploration of self-compassion and self-criticism in the context of personal recovery from borderline personality disorder. *Australasian Psychiatry*, 27, 56–59.
- Dore, J., Turnipseed, B., Dwyer, S., Turnipseed, A., Andries, J., Ascani, G., & Wolfson, P. (2019). Ketamine assisted psychotherapy (KAP): Patient demographics, clinical data and outcomes in three large practices administering ketamine with psychotherapy. *Journal of Psychoactive Drugs*, 51, 189–198. <https://doi.org/10.1080/02791072.2019.1587556>.
- Elices, M., Pascual, J. C., Portella, M. J., Feliu-Soler, A., Martín-Blanco, A., Carmona, C., & Soler, J. (2016). Impact of mindfulness training on borderline personality disorder: A randomized trial. *Mindfulness*, 7, 584–595.
- Erritzoe, D., Roseman, L., Nour, M. M., MacLean, K., Kaelen, M., Nutt, D. J., & Carhart-Harris, R. L. (2018). Effects of psilocybin therapy on personality structure. *Acta Psychiatrica Scandinavica*, 138, 368–378.
- Feifel, D., Malcolm, B., Boggie, D., & Lee, K. (2017). Low-dose ketamine for treatment resistant depression in an academic clinical practice setting. *Journal of Affective Disorders*, 221, 283–288.
- Feliu-Soler, A., Pascual, J. C., Borrás, X., Portella, M. J., Martín-Blanco, A., Armario, A., ... Soler, J. (2014). Effects of dialectical behaviour therapy-mindfulness training on emotional reactivity in borderline personality disorder: Preliminary results. *Clinical Psychology & Psychotherapy*, 21, 363–370.
- Feliu-Soler, A., Pascual, J. C., Elices, M., Martín-Blanco, A., Carmona, C., Cebolla, A., ... Soler, J. (2017). Fostering self-compassion and loving-kindness in patients with borderline personality disorder: A randomized pilot study. *Clinical Psychology &*

- Psychotherapy*, 24, 278–286.
- Fernández, X., dos Santos, R. G., Cutchet, M., Fondevila, S., González, D., Alcázar, M.Á., ... Fábregas, J. M. (2014). Assessment of the psychotherapeutic effects of ritual ayahuasca use on drug dependency: A pilot study. In B. C. Labate, & C. Cavnar (Eds.). *The therapeutic use of ayahuasca* (pp. 183–196). New York, NY: Springer Heidelberg.
- Francois, D., Roth, S. D., & Klingman, D. (2015). The efficacy of pharmacotherapy for borderline personality disorder: A review of the available randomized controlled trials. *Psychiatric Annals*, 45, 431–437.
- Franquesa, A., Sainz-Cort, A., Gandy, S., Soler, J., Alcázar-Córcoles, M.Á., & Bouso, J. C. (2018). Psychological variables implied in the therapeutic effect of ayahuasca: A contextual approach. *Psychiatry Research*, 264, 334–339.
- Frye, C. G., Wardle, M. C., Norman, G. J., & de Wit, H. (2014). MDMA decreases the effects of simulated social rejection. *Pharmacology Biochemistry and Behavior*, 117, 1–6.
- Gabay, A. S., Kempton, M. J., Gillean, J., & Mehta, M. A. (2019). MDMA increases co-operation and recruitment of social brain areas when playing trustworthy players in an iterated Prisoner's Dilemma. *Journal of Neuroscience*, 39, 307–320.
- Gad, M. A., Pucker, H. E., Hein, K. E., Temes, C. M., Frankenburg, F. R., Fitzmaurice, G. M., & Zanarini, M. C. (2019). Facets of identity disturbance reported by patients with borderline personality disorder and personality-disordered comparison subjects over 20 years of prospective follow-up. *Psychiatry Research*, 271, 76–82.
- Gasser, P. (1994). Psychedelic therapy with MDMA and LSD in Switzerland. *MAPS Newsletter*, 5, 3–7.
- Gasser, P., Kirchner, K., & Passie, T. (2015). LSD-assisted psychotherapy for anxiety associated with a life-threatening disease: A qualitative study of acute and sustained subjective effects. *Journal of Psychopharmacology*, 29, 57–68.
- Goldsmith, N. M. (2007). The ten lessons of psychedelic psychotherapy, rediscovered. *Psychiatric medicine: New evidence for hallucinogenic substances as treatments*, 2, 107–141.
- Gratz, K. L., Bardeen, J. R., Levy, R., Dixon-Gordon, K. L., & Tull, M. T. (2015). Mechanisms of change in an emotion regulation group therapy for deliberate self-harm among women with borderline personality disorder. *Behaviour Research and Therapy*, 65, 29–35.
- Gratz, K. L., Levy, R., & Tull, M. T. (2012). Emotion regulation as a mechanism of change in an acceptance-based emotion regulation group therapy for deliberate self-harm among women with borderline personality pathology. *Journal of Cognitive Psychotherapy*, 26, 365–380.
- Gregory, W. E., Glazer, J. V., & Berenson, K. R. (2017). Self-compassion, self-injury, and pain. *Cognitive Therapy and Research*, 41, 777–786.
- Griffiths, R. R., Johnson, M. W., Carducci, M. A., Umbricht, A., Richards, W. A., Richards, B. D., ... Klinedinst, M. A. (2016). Psilocybin produces substantial and sustained decreases in depression and anxiety in patients with life-threatening cancer: A randomized double-blind trial. *Journal of Psychopharmacology*, 30, 1181–1197.
- Griffiths, R. R., Johnson, M. W., Richards, W. A., Richards, B. D., Jesse, R., MacLean, K. A., ... Klinedinst, M. A. (2018). Psilocybin-occasioned mystical-type experience in combination with meditation and other spiritual practices produces enduring positive changes in psychological functioning and in trait measures of prosocial attitudes and behaviors. *Journal of Psychopharmacology*, 32, 49–69.
- Griffiths, R. R., Johnson, M. W., Richards, W. A., Richards, B. D., McCann, U., & Jesse, R. (2011). Psilocybin occasioned mystical-type experiences: Immediate and persisting dose-related effects. *Psychopharmacology*, 218, 649–665.
- Griffiths, R. R., Richards, W. A., Johnson, M. W., McCann, U. D., & Jesse, R. (2008). Mystical-type experiences occasioned by psilocybin mediate the attribution of personal meaning and spiritual significance 14 months later. *Journal of Psychopharmacology*, 22, 621–632.
- Griffiths, R. R., Richards, W. A., McCann, U., & Jesse, R. (2006). Psilocybin can occasion mystical-type experiences having substantial and sustained personal meaning and spiritual significance. *Psychopharmacology*, 187, 268–283.
- Grunebaum, M. F., Galfalvy, H. C., Choo, T. H., Keilp, J. G., Moitra, V. K., Parris, M. S., ... Oquendo, M. A. (2017). Ketamine for rapid reduction of suicidal thoughts in major depression: A midazolam-controlled randomized clinical trial. *American Journal of Psychiatry*, 175, 327–335.
- Gunderson, J. G., Stout, R. L., McGlashan, T. H., Shea, M. T., Morey, L. C., Grilo, C. M., ... Ansell, E. (2011). Ten-year course of borderline personality disorder: Psychopathology and function from the Collaborative Longitudinal Personality Disorders study. *Archives of General Psychiatry*, 68, 827–837.
- Hamill, J., Hallak, J., Dursun, S. M., & Baker, G. (2019). Ayahuasca: Psychological and physiologic effects, pharmacology and potential uses in addiction and mental illness. *Current Neuropharmacology*, 17, 108–128.
- Hancock-Johnson, E., Griffiths, C., & Picchioni, M. (2017). A focused systematic review of pharmacological treatment for borderline personality disorder. *CNS Drugs*, 31, 345–356.
- Hasking, P., Boyes, M. E., Finlay-Jones, A., McEvoy, P. M., & Rees, C. S. (2019). Common pathways to NSSI and suicide ideation: The roles of rumination and self-compassion. *Archives of Suicide Research*, 23, 247–260.
- Heath, L. M., Laporte, L., Paris, J., Hamdullahpur, K., & Gill, K. J. (2018). Substance misuse is associated with increased psychiatric severity among treatment-seeking individuals with borderline personality disorder. *Journal of Personality Disorders*, 32, 694–708.
- Hendricks, P. S., Crawford, M. S., Cropsey, K. L., Copes, H., Sweat, N. W., Walsh, Z., & Pavela, G. (2017). The relationships of classic psychedelic use with criminal behavior in the United States adult population. *Journal of Psychopharmacology*, 32, 37–48.
- Hendricks, P. S., Thorne, C. B., Clark, C. B., Coombs, D. W., & Johnson, M. W. (2015). Classic psychedelic use is associated with reduced psychological distress and suicidality in the United States adult population. *Journal of Psychopharmacology*, 29, 280–288.
- Herpertz, S. C., Schneider, I., Schmahl, C., & Bertsch, K. (2018). Neurobiological mechanisms mediating emotion dysregulation as targets of change in borderline personality disorder. *Psychopathology*, 51, 96–104.
- Hysek, C. M., Schmid, Y., Simmler, L. D., Domes, G., Heinrichs, M., Eisenegger, C., ... Liechti, M. E. (2013). MDMA enhances emotional empathy and prosocial behavior. *Social Cognitive and Affective Neuroscience*, 9, 1645–1652.
- Johnson, M. W., Garcia-Romeu, A., Cosimano, M. P., & Griffiths, R. R. (2014). Pilot study of the 5-HT_{2A} agonist psilocybin in the treatment of tobacco addiction. *Journal of Psychopharmacology*, 28, 983–992.
- Johnson, M. W., Griffiths, R. R., Hendricks, P. S., & Henningfield, J. E. (2018). The abuse potential of medical psilocybin according to the 8 factors of the Controlled Substances Act. *Neuropharmacology*, 142, 143–166.
- Johnson, M. W., Hendricks, P. S., Barrett, F. S., & Griffiths, R. R. (2019). Classic psychedelics: An integrative review of epidemiology, mystical experience, brain network function, and therapeutics. *Pharmacology & Therapeutics*, 197, 83–102. <https://doi.org/10.1016/j.pharmthera.2018.11.010>.
- Johnson, M. W., Richards, W. A., & Griffiths, R. R. (2008). Human hallucinogen research: Guidelines for safety. *Journal of Psychopharmacology*, 22, 603–620.
- Jørgensen, C. R., Berntsen, D., Bech, M., Kjølbjerg, M., Bennedsen, B. E., & Ramsgaard, S. B. (2012). Identity-related autobiographical memories and cultural life scripts in patients with borderline personality disorder. *Consciousness and Cognition*, 21, 788–798.
- Kabat-Zinn, J. (1990). *Full catastrophe living: Using the wisdom of your body and mind to face stress, pain, and illness*. New York, NY: Dell Publishing.
- Kaelin, M., Giribaldi, B., Raine, J., Evans, L., Timmerman, C., Rodriguez, N., & Carhart-Harris, R. (2018). The hidden therapist: Evidence for a central role of music in psychedelic therapy. *Psychopharmacology*, 235, 505–519.
- Kamboj, S. K., Kilford, E. J., Minchin, S., Moss, A., Lawn, W., Das, R. K., ... Freeman, T. P. (2015). Recreational 3, 4-methylenediox-N-methylamphetamine (MDMA) or 'ecstasy' and self-focused compassion: Preliminary steps in the development of a therapeutic psychopharmacology of contemplative practices. *Journal of Psychopharmacology*, 29, 961–970.
- Kamboj, S. K., Wallden, Y. S., Falconer, C. J., Alotaibi, M. R., Blagbrough, I. S., Husbands, S. M., & Freeman, T. P. (2017). Additive effects of 3, 4-methylenediox-N-methylamphetamine (MDMA) and compassionate imagery on self-compassion in recreational users of ecstasy. *Mindfulness*, 9, 1134–1145.
- Katsakou, C., & Pistrang, N. (2017). Clients' experiences of treatment and recovery in borderline personality disorder: A meta-synthesis of qualitative studies. *Psychotherapy Research*, 28, 940–957.
- Katsakou, C., Pistrang, N., Barnicot, K., White, H., & Priebe, S. (2017). Processes of recovery through routine or specialist treatment for borderline personality disorder (BPD): A qualitative study. *Journal of Mental Health*. <https://doi.org/10.1080/09638237.2017.1340631>.
- Kazdin, A. E. (2009). Understanding how and why psychotherapy leads to change. *Psychotherapy Research*, 19, 418–428.
- Kernberg, O. F. (1975). *Borderline conditions and pathological narcissism*. New York, NY: Jason Aronson.
- Kerr, C. E., Josyula, K., & Littenberg, R. (2011). Developing an observing attitude: An analysis of meditation diaries in an MBSR clinical trial. *Clinical Psychology and Psychotherapy*, 18, 80–93.
- Kim, J., & Mierzwinski-Urban, M. (2017). *Ketamine for treatment-resistant depression or post-traumatic stress disorder in various settings: A review of clinical effectiveness, safety, and guidelines*. Ottawa, ON: Canadian Agency for Drugs and Technologies in Health.
- Kometer, M., Schmidt, A., Bachmann, R., Studerus, E., Seifritz, E., & Vollenweider, F. X. (2012). Psilocybin biases facial recognition, goal-directed behavior, and mood state toward positive relative to negative emotions through different serotonergic sub-receptors. *Biological Psychiatry*, 72, 898–906.
- Kopala-Sibley, D. C., Zuroff, D. C., Russell, J. J., Moskowitz, D. S., & Paris, J. (2012). Understanding heterogeneity in borderline personality disorder: Differences in affective reactivity explained by the traits of dependency and self-criticism. *Journal of Abnormal Psychology*, 121, 680–691.
- Kostowski, W., Rewerski, W., & Piechocki, T. (1972). The effects of some hallucinogens on aggressiveness of mice and rats. *Pharmacology*, 7, 259–263.
- Kraehenmann, R., Schmidt, A., Friston, K., Preller, K. H., Seifritz, E., & Vollenweider, F. X. (2016). The mixed serotonin receptor agonist psilocybin reduces threat-induced modulation of amygdala connectivity. *NeuroImage: Clinical*, 11, 53–60.
- Krantz, L. H., McMain, S., & Kuo, J. R. (2018). The unique contribution of acceptance without judgment in predicting nonsuicidal self-injury after 20-weeks of dialectical behaviour therapy group skills training. *Behaviour Research and Therapy*, 104, 44–50.
- Krawitz, R. (2012). Behavioural treatment of chronic, severe self-loathing in people with borderline personality disorder. Part 1: Interrupting the self-loathing cycle. *Australasian Psychiatry*, 20, 419–424.
- Krebs, T. S., & Johansen, P. Ø. (2012). Lysergic acid diethylamide (LSD) for alcoholism: Meta-analysis of randomized controlled trials. *Journal of Psychopharmacology*, 26, 994–1002.
- Krupitsky, E. M., & Grinenko, A. Y. (1996). Ketamine psychotherapy: Results and mechanisms. *International Journal of Transpersonal Studies*, 15, 32–66.
- Lafrance, A., Loizaga-Velder, A., Fletcher, J., Renelli, M., Files, N., & Tupper, K. W. (2017). Nourishing the spirit: Exploratory research on ayahuasca experiences along the continuum of recovery from eating disorders. *Journal of Psychoactive Drugs*, 49, 427–435.
- Lauritsen, C., Mazuera, S., Lipton, R. B., & Ashina, S. (2016). Intravenous ketamine for subacute treatment of refractory chronic migraine: A case series. *The Journal of Headache and Pain*, 17, 106. <https://doi.org/10.1186/s10194-016-0700-3>.
- Lazarus, S. A., Cheavens, J. S., Festa, F., & Rosenthal, M. Z. (2014). Interpersonal functioning in borderline personality disorder: A systematic review of behavioral and laboratory-based assessments. *Clinical Psychology Review*, 34, 193–205.

- Lieb, K., Völm, B., Rücker, G., Timmer, A., & Stoffers, J. M. (2010). Pharmacotherapy for borderline personality disorder: Cochrane systematic review of randomised trials. *The British Journal of Psychiatry*, 196, 4–12.
- Linehan, M. (1993). *Cognitive-behavioral treatment of borderline personality disorder*. New York, NY: Guilford Press.
- Linehan, M. M., Comtois, K. A., Murray, A. M., Brown, M. Z., Gallop, R. J., Heard, H. L., & Lindenboim, N. (2006). Two-year randomized controlled trial and follow-up of dialectical behavior therapy vs therapy by experts for suicidal behaviors and borderline personality disorder. *Archives of General Psychiatry*, 63, 757–766.
- Linehan, M. M., McDavid, J. D., Brown, M. Z., Sayrs, J. H., & Gallop, R. J. (2008). Olanzapine plus dialectical behavior therapy for women with high irritability who meet criteria for borderline personality disorder: A double-blind, placebo-controlled pilot study. *Journal of Clinical Psychiatry*, 69, 999–1005.
- Ly, C., Greb, A. C., Cameron, L. P., Wong, J. M., Barragan, E. V., Wilson, P. C., ... Duim, W. C. (2018). Psychedelics promote structural and functional neural plasticity. *Cell Reports*, 23, 3170–3182.
- MacLean, K. A., Johnson, M. W., & Griffiths, R. R. (2011). Mystical experiences occasioned by the hallucinogen psilocybin lead to increases in the personality domain of openness. *Journal of Psychopharmacology*, 25, 1453–1461.
- Malone, T. C., Mennenga, S. E., Guss, J., Podrebarac, S. K., Owens, L. T., Bossis, A. P., ... Ross, S. (2018). Individual experiences in four cancer patients following psilocybin-assisted psychotherapy. *Frontiers in Pharmacology*, 9, 256. <https://doi.org/10.3389/fphar.2018.00256>.
- Marin, L. K., Kapil-Par, K. N., Harris, R. E., & Goodman, M. (2018). Combination treatments in borderline personality disorder: Bridging the gap between clinical practice and empirical data. *Current Treatment Options in Psychiatry*, 5, 141–161.
- McCabe, O. L., Savage, C., Kurland, A., & Unger, S. (1972). Psychedelic (LSD) therapy of neurotic disorders: Short term effects. *Journal of Psychedelic Drugs*, 5, 18–28.
- McMain, S., Links, P. S., Guimond, T., Wnuk, S., Eynan, R., Bergmans, Y., & Warwar, S. (2013). An exploratory study of the relationship between changes in emotion and cognitive processes and treatment outcome in borderline personality disorder. *Psychotherapy Research*, 23, 658–673.
- Mitchell, R., Roberts, R., Bartsch, D., & Sullivan, T. (2019). Changes in mindfulness facets in a dialectical behaviour therapy skills training group program for borderline personality disorder. *Journal of Clinical Psychology*, 75, 958–969. <https://doi.org/10.1002/jclp.22744>.
- Mithoefer, M. C., Wagner, M. T., Mithoefer, A. T., Jerome, L., & Doblin, R. (2011). The safety and efficacy of \pm 3, 4-methylenedioxymethamphetamine-assisted psychotherapy in subjects with chronic, treatment-resistant posttraumatic stress disorder: The first randomized controlled pilot study. *Journal of Psychopharmacology*, 25, 439–452.
- Mithoefer, M. C., Wagner, M. T., Mithoefer, A. T., Jerome, L., Martin, S. F., Yazar-Klosinski, B., ... Doblin, R. (2013). Durability of improvement in post-traumatic stress disorder symptoms and absence of harmful effects or drug dependency after 3, 4-methylenedioxymethamphetamine-assisted psychotherapy: A prospective long-term follow-up study. *Journal of Psychopharmacology*, 27, 28–39.
- Molero, P., Ramos-Quiroga, J. A., Martín-Santos, R., Calvo-Sánchez, E., Gutiérrez-Rojas, L., & Meana, J. J. (2018). Antidepressant efficacy and tolerability of ketamine and esketamine: A critical review. *CNS Drugs*, 32, 411–420.
- Multidisciplinary Association for Psychedelics Studies. (2017). MDMA investigational brochure. Retrieved from <https://maps.org/research/mdma/mdma-research-timeline/104-other-mdma-resources/5400-mdma-investigator-s-brochure-and-fda-annual-reportanditemid=485>.
- Murrough, J. W., Soleimani, L., DeWilde, K. E., Collins, K. A., Lapidus, K. A., Iacoviello, B. M., & Price, R. B. (2015). Ketamine for rapid reduction of suicidal ideation: A randomized controlled trial. *Psychological Medicine*, 45, 3571–3580.
- National Institute of Health and Clinical Excellence. (2009). *Borderline personality disorder: Treatment and management*. London, UK: Author.
- Newhill, C. E., Eack, S. M., & Mulvey, E. P. (2009). Violent behavior in borderline personality. *Journal of Personality Disorders*, 23, 541–554.
- Nicholas, C. R., Henriquez, K. M., Gassman, M. C., Cooper, K. M., Muller, D., Hetzel, S., ... Hutson, P. R. (2018). High dose psilocybin is associated with positive subjective effects in healthy volunteers. *Journal of Psychopharmacology*, 32, 770–778.
- Nichols, D. E. (2004). Hallucinogens. *Pharmacology and Therapeutics*, 101, 131–181.
- Nichols, D. E. (2016). Psychedelics. *Pharmacological Reviews*, 68, 264–355.
- Nichols, C. D., & Sanders-Bush, E. (2002). A single dose of lysergic acid diethylamide influences gene expression patterns within the mammalian brain. *Neuropsychopharmacology*, 26, 634–642.
- Nour, M. M., & Carhart-Harris, R. L. (2017). Psychedelics and the science of self-experience. *The British Journal of Psychiatry*, 210, 177–179.
- Nutt, D. J., King, L. A., & Nichols, D. E. (2013). Effects of Schedule I drug laws on neuroscience research and treatment innovation. *Nature Reviews Neuroscience*, 14, 577–585.
- Nutt, D. J., King, L. A., & Phillips, L. D. (2010). Drug harms in the UK: A multicriteria decision analysis. *The Lancet*, 376, 1558–1565.
- Orzech, K. M., Shapiro, S. L., Brown, K. W., & McKay, M. (2009). Intensive mindfulness training-related changes in cognitive and emotional experience. *The Journal of Positive Psychology*, 4, 212–222.
- Ot'ala, G. M., Grigsby, J., Poulter, B., Van Derveer, J. W., III, Giron, S. G., Jerome, L., ... Mithoefer, M. C. (2018). 3, 4-methylenedioxymethamphetamine-assisted psychotherapy for treatment of chronic posttraumatic stress disorder: A randomized phase 2 controlled trial. *Journal of Psychopharmacology*, 32, 1295–1307.
- O'Toole, S. K., Diddy, E., & Kent, M. (2012). Mindfulness and emotional well-being in women with borderline personality disorder. *Mindfulness*, 3, 117–123.
- Palhano-Fontes, F., Barreto, D., Onias, H., Andrade, K. C., Novais, M. M., Pessoa, J. A., ... Tófoli, L. F. (2019). Rapid antidepressant effects of the psychedelic ayahuasca in treatment-resistant depression: A randomized placebo-controlled trial. *Psychological Medicine*, 49, 655–663.
- Paris, J. (2009). The treatment of borderline personality disorder: Implications of research on diagnosis, etiology, and outcome. *Annual Review of Clinical Psychology*, 5, 277–290.
- Paton, C., Crawford, M. J., Bhatti, S. F., Patel, M. X., & Barnes, T. R. (2015). The use of psychotropic medication in patients with emotionally unstable personality disorder under the care of UK mental health services. *Journal of Clinical Psychiatry*, 76, 512–518.
- Perroud, N., Nicastro, R., Jermann, F., & Huguelet, P. (2012). Mindfulness skills in borderline personality disorder patients during dialectical behavior therapy: Preliminary results. *International Journal of Psychiatry in Clinical Practice*, 16, 189–196.
- Pisano, V. D., Putnam, N. P., Kramer, H. M., Franciotti, K. J., Halpern, J. H., & Holden, S. C. (2017). The association of psychedelic use and opioid use disorders among illicit users in the United States. *Journal of Psychopharmacology*, 31, 606–613.
- Pokorny, T., Preller, K. H., Kometer, M., Dziobek, I., & Vollenweider, F. X. (2017). Effect of psilocybin on empathy and moral decision-making. *International Journal of Neuropsychopharmacology*, 20, 747–757.
- Preller, K. H., Pokorny, T., Hock, A., Kraehenmann, R., Stämpfli, P., Seifritz, E., ... Vollenweider, F. X. (2016). Effects of serotonin 2A/1A receptor stimulation on social exclusion processing. *Proceedings of the National Academy of Sciences*, 113, 5119–5124.
- Preller, K. H., Pokorny, T., Krähenmann, R., Scheidegger, M., Dziobek, I., Stämpfli, P., & Vollenweider, F. X. (2015). The 5-HT_{2A/1A} agonist psilocybin reduces social pain and enhances empathy in healthy volunteers. *European Neuropsychopharmacology*, 25, S301.
- Renneberg, B., Herm, K., Hahn, A., Staebler, K., Lammers, C. H., & Roepke, S. (2012). Perception of social participation in borderline personality disorder. *Clinical Psychology & Psychotherapy*, 19, 473–480.
- Rizvi, S. L., & Thomas, M. C. (2017). Borderline personality disorder: Underlying mechanisms and implications for treatment. In D. McKay, J. S. Abramowitz, & E. A. Storch (Eds.). *Treatments for psychological problems and syndromes* (pp. 360–375). Hoboken, NJ: John Wiley & Sons.
- Ross, S., Bossis, A., Guss, J., Agin-Lieb, G., Malone, T., Cohen, B., ... Su, Z. (2016). Rapid and sustained symptom reduction following psilocybin treatment for anxiety and depression in patients with life-threatening cancer: A randomized controlled trial. *Journal of Psychopharmacology*, 30, 1165–1180.
- Rudge, S., Feigenbaum, J. D., & Fonagy, P. (2017). Mechanisms of change in dialectical behaviour therapy and cognitive behaviour therapy for borderline personality disorder: A critical review of the literature. *Journal of Mental Health*. <https://doi.org/10.1080/09638237.2017.1322185>.
- Sampedro, F., de la Fuente Revenga, M., Valle, M., Roberto, N., Domínguez-Clavé, E., Elices, M., ... Friedlander, P. (2017). Assessing the psychedelic “after-glow” in ayahuasca users: Post-acute neurometabolic and functional connectivity changes are associated with enhanced mindfulness capacities. *International Journal of Neuropsychopharmacology*, 20, 698–711.
- Sanches, R. F., de Lima Osório, F., Dos Santos, R. G., Macedo, L. R., Maia-de-Oliveira, J. P., Wichert-Ana, L., ... Hallak, J. E. (2016). Antidepressant effects of a single dose of ayahuasca in patients with recurrent depression: A spect study. *Journal of Clinical Psychopharmacology*, 36, 77–81.
- Sansone, R. A., & Wiederman, M. W. (2009). The abuse of prescription medications: Borderline personality patients in psychiatric versus non-psychiatric settings. *The International Journal of Psychiatry in Medicine*, 39, 147–154.
- dos Santos, R. G., Osorio, F. L., Crippa, J. A. S., & Hallak, J. E. (2016). Classical hallucinogens and neuroimaging: A systematic review of human studies: Hallucinogens and neuroimaging. *Neuroscience & Biobehavioral Reviews*, 71, 715–728.
- Schmid, Y., & Liechti, M. E. (2018). Long-lasting subjective effects of LSD in normal subjects. *Psychopharmacology*, 235, 535–545.
- Schulze, L., Schmal, C., & Niedtfeld, I. (2016). Neural correlates of disturbed emotion processing in borderline personality disorder: A multimodal meta-analysis. *Biological Psychiatry*, 79, 97–106.
- Sebastian, A., Jacob, G., Lieb, K., & Tüscher, O. (2013). Impulsivity in borderline personality disorder: A matter of disturbed impulse control or a facet of emotional dysregulation? *Current Psychiatry Reports*, 15, 339.
- Sessa, B., Higbed, L., & Nutt, D. (2019). A review of 3, 4-methylenedioxymethamphetamine (MDMA)-assisted psychotherapy. *Frontiers in Psychiatry*, 10. <https://doi.org/10.3389/fpsy.2019.00138>.
- Smigielski, L., Scheidegger, M., Kometer, M., & Vollenweider, F. X. (2019). Psilocybin-assisted mindfulness training modulates self-consciousness and brain default mode network connectivity with lasting effects. *NeuroImage*, 196, 207–215.
- Soler, J., Elices, M., Domínguez-Clavé, E., Pascual, J. C., Feilding, A., Navarro-Gil, M., & Riba, J. (2018). Four weekly ayahuasca sessions lead to increases in “acceptance” capacities: A comparison study with a standard 8-week mindfulness training program. *Frontiers in Pharmacology*, 9, 224.
- Soler, J., Elices, M., Franquesa, A., Barker, S., Friedlander, P., Feilding, A., ... Riba, J. (2016). Exploring the therapeutic potential of Ayahuasca: Acute intake increases mindfulness-related capacities. *Psychopharmacology*, 233, 823–829.
- Soler, J., Valdepérez, A., Feliu-Soler, A., Pascual, J. C., Portella, M. J., Martín-Blanco, A., ... Pérez, V. (2012). Effects of the dialectical behavior therapy-mindfulness module on attention in patients with borderline personality disorder. *Behaviour Research and Therapy*, 50, 150–157.
- Soloff, P. H., Lis, J. A., Kelly, T., Cornelius, J., & Ulrich, R. (1994). Self-mutilation and suicidal behavior in borderline personality disorder. *Journal of Personality Disorders*, 8, 257–267.
- Soloff, P. H., Lynch, K. G., & Kelly, T. M. (2002). Childhood abuse as a risk factor for suicidal behavior in borderline personality disorder. *Journal of Personality Disorders*,

- 16, 201–214.
- Staebler, K., Helbing, E., Rosenbach, C., & Renneberg, B. (2011). Rejection sensitivity and borderline personality disorder. *Clinical Psychology & Psychotherapy*, 18, 275–283.
- Staebler, K., Renneberg, B., Stopsack, M., Fiedler, P., Weiler, M., & Roepke, S. (2011). Facial emotional expression in reaction to social exclusion in borderline personality disorder. *Psychological Medicine*, 41, 1929–1938.
- Starcevic, V., & Janca, A. (2018). Pharmacotherapy of borderline personality disorder: Replacing confusion with prudent pragmatism. *Current Opinion in Psychiatry*, 31, 69–73.
- Stepp, S. D., Epler, A. J., Jahng, S., & Trull, T. J. (2008). The effect of dialectical behavior therapy skills use on borderline personality disorder features. *Journal of Personality Disorders*, 22, 549–563.
- Stoffers, J. M., & Lieb, K. (2015). Pharmacotherapy for borderline personality disorder—current evidence and recent trends. *Current Psychiatry Reports*, 17, 534.
- Stoffers, J. M., Völlm, B. A., Rücker, G., Timmer, A., Huband, N., & Lieb, K. (2012). Psychological therapies for people with borderline personality disorder. *Cochrane Database of Systematic Reviews*, 1–186. <https://doi.org/10.1002/14651858.CD005652>.
- Stratton, N., Alvarez, M. M., Labrish, C., Barnhart, R., & McMain, S. (2018). Predictors of dropout from a 20-week dialectical behavior therapy skills group for suicidal behaviors and borderline personality disorder. *Journal of Personality Disorders*. <https://doi.org/10.1521/pedi.2018.32.391>.
- Strong, C. E., & Kabbaj, M. (2018). On the safety of repeated ketamine infusions for the treatment of depression: Effects of sex and developmental periods. *Neurobiology of Stress*, 9, 166–175.
- Stroud, J. B., Freeman, T. P., Leech, R., Hindocha, C., Lawn, W., Nutt, D. J., ... Carhart-Harris, R. L. (2018). Psilocybin with psychological support improves emotional face recognition in treatment-resistant depression. *Psychopharmacology*, 235, 459–466.
- Temes, C. M., Frankenburg, F. R., Fitzmaurice, G. M., & Zanarini, M. C. (2019). Deaths by suicide and other causes among patients with borderline personality disorder and personality-disordered comparison subjects over 24 years of prospective follow-up. *The Journal of Clinical Psychiatry*, 80. <https://doi.org/10.4088/JCP.18m12436>.
- Terzi, L., Martino, F., Berardi, D., Bortolotti, B., Sasdelli, A., & Menchetti, M. (2017). Aggressive behavior and self-harm in Borderline Personality Disorder: The role of impulsivity and emotion dysregulation in a sample of outpatients. *Psychiatry Research*, 249, 321–326.
- Thiessen, M. S., Walsh, Z., Bird, B. M., & Lafrance, A. (2018). Psychedelic use and intimate partner violence: The role of emotion regulation. *Journal of Psychopharmacology*, 32, 749–755.
- Thomas, G., Lucas, P., Capler, N. R., Tupper, K. W., & Martin, G. (2013). Ayahuasca-assisted therapy for addiction: Results from a preliminary observational study in Canada. *Current Drug Abuse Reviews*, 6, 30–42.
- Tomlinson, M. F., Brown, M., & Hoaken, P. N. (2016). Recreational drug use and human aggressive behavior: A comprehensive review since 2003. *Aggression and Violent Behavior*, 27, 9–29.
- Torgersen, S. (2012). Epidemiology. In T. A. Widiger (Ed.). *The Oxford handbook of personality disorders* (pp. 186–205). New York, NY: Oxford University Press.
- Tortella-Feliu, M., Soler, J., Burns, L., Cebolla, A., Elices, M., Pascual, J. C., & García-Campayo, J. (2018). Relationship between effortful control and facets of mindfulness in meditators, non-meditators and individuals with borderline personality disorder. *Personality and Mental Health*, 12, 265–278.
- Trull, T. J., Solhan, M. B., Brown, W. C., Tomko, R. L., Schaefer, L., McLaughlin, K. D., & Jahng, S. (2016). Substance use disorders and personality disorders. In K. Sher (Ed.). *The Oxford handbook of substance use disorders*. New York, NY: Oxford University Press.
- Van Asselt, A. D. I., Dirksen, C. D., Arntz, A., & Severens, J. L. (2007). The cost of borderline personality disorder: Societal cost of illness in BPD-patients. *European Psychiatry*, 22, 354–361.
- Vollenweider, F. X., & Kometer, M. (2010). The neurobiology of psychedelic drugs: Implications for the treatment of mood disorders. *Nature Reviews Neuroscience*, 11, 642–651.
- Wagner, A. C., Mithoefer, M. C., Mithoefer, A. T., & Monson, C. M. (2019). Combining cognitive-behavioral conjoint therapy for PTSD with 3, 4-methylenedioxymethamphetamine (MDMA): A case example. *Journal of Psychoactive Drugs*, 51, 166–173. <https://doi.org/10.1080/02791072.2019.1589028>.
- Wagner, M. T., Mithoefer, M. C., Mithoefer, A. T., MacAulay, R. K., Jerome, L., Yazar-Klosinski, B., & Doblin, R. (2017). Therapeutic effect of increased openness: Investigating mechanism of action in MDMA-assisted psychotherapy. *Journal of Psychopharmacology*, 31, 967–974.
- Walsh, Z., Hendricks, P. S., Smith, S., Kosson, D. S., Thiessen, M. S., Lucas, P., & Swogger, M. T. (2016). Hallucinogen use and intimate partner violence: Prospective evidence consistent with protective effects among men with histories of problematic substance use. *Journal of Psychopharmacology*, 30, 601–607.
- Walsh, Z., & Thiessen, M. S. (2018). Psychedelics and the new behaviourism: Considering the integration of third-wave behaviour therapies with psychedelic-assisted therapy. *International Review of Psychiatry*, 30, 343–349.
- Watts, R., Day, C., Krzanowski, J., Nutt, D., & Carhart-Harris, R. (2017). Patients' accounts of increased “connectedness” and “acceptance” after psilocybin for treatment-resistant depression. *Journal of Humanistic Psychology*, 57, 520–564.
- Wilson, S.T., Fertuck, E. A., Kwitel, A., Stanley, M. C., & Stanley, B. (2006). Impulsivity, suicidality and alcohol use disorders in adolescents and young adults with borderline personality disorder. *International Journal of Adolescent Medicine and Health*, 18, 189–196.
- Winter, D., Bohus, M., & Lis, S. (2017). Understanding negative self-evaluations in borderline personality disorder—A review of self-related cognitions, emotions, and motives. *Current Psychiatry Reports*, 19, 1–9.
- Wnuk, S., McMain, S., Links, P. S., Habinski, L., Murray, J., & Guimond, T. (2013). Factors related to dropout from treatment in two outpatient treatments for borderline personality disorder. *Journal of Personality Disorders*, 27, 716–726.
- Wupperman, P., Fickling, M., Klemanski, D. H., Berking, M., & Whitman, J. B. (2013). Borderline personality features and harmful dysregulated behavior: The mediational effect of mindfulness. *Journal of Clinical Psychology*, 69, 903–911.
- Zanarini, M. C., Frankenburg, F. R., Hennen, J., Reich, D. B., & Silk, K. R. (2004). Axis I comorbidity in patients with borderline personality disorder: 6-year follow-up and prediction of time to remission. *American Journal of Psychiatry*, 161, 2108–2114.
- Zanarini, M. C., Frankenburg, F. R., Hennen, J., & Silk, K. R. (2004). Mental health service utilization by borderline personality disorder patients and Axis II comparison subjects followed prospectively for 6 years. *Journal of Clinical Psychiatry*, 65, 28–36.
- Zanarini, M. C., Frankenburg, F. R., Reich, D. B., Silk, K. R., Hudson, J. I., & McSweeney, L. B. (2007). The subsyndromal phenomenology of borderline personality disorder: A 10-year follow-up study. *American Journal of Psychiatry*, 164, 929–935.
- Zanarini, M. C., Temes, C. M., Frankenburg, F. R., Reich, D. B., & Fitzmaurice, G. M. (2019). Levels of acceptance and forgiveness reported by patients with BPD and personality-disordered comparison subjects over 20 years of prospective follow-up. *Journal of Personality Disorders*. <https://doi.org/10.1521/pedi.2019.33.395>.
- Zeifman, R. J., Boritz, T. Z., Barnhart, R., Labrish, C., & McMain, S. F. (2019). The independent roles of mindfulness and distress tolerance in treatment outcomes in dialectical behavior therapy skills training. *Personality Disorders: Theory, Research, and Treatment*. <https://doi.org/10.1037/per0000368> Advance online publication.